



AIUB DSpace Publication Details

Title: A study on CNT/Peptide interface by second harmonic generation (SHG) method

Author(s) Name: Md. Ehasanul Haque, Md. Masud Parvez, Humayra Ferdous, Md. Abdus Sattar, Md. Mizanur Rahman and Md. Asadul Hoque

Contact Email(s): ehasanul@aiub.edu

Published Conference Name: International Conference on Physics 2020

Type of Publication: International Conference

Volume: _____ **Issue:** _____

Publisher: _____

Publication Date: March 2020

ISSN: _____

DOI: _____

URL: _____

Other Related Info.: _____



AIUB DSpace Publication Details

Abstract:

We observed the second harmonic generation (SHG) intensity from the peptide molecules absorbed carbon nanotube surface grown on the Si/SiO₂/Co substrate. We dropped different concentrations of peptide molecules such as 100nM, 1μM and 10μM on the three Si/SiO₂/Co/CNT substrate individually. The SHG intensity was measured from the CNT/PEP interface by using 1.17 eV pulsed laser light. The results show that, the SHG intensity increased with increasing the peptide concentrations. In order to confirm about the SHG signal detected from the CNT/PEP interface, we dropped similar concentrations of peptide molecule on the three different Si/SiO₂ substrate having no CNT layer individually and we observed the same results. In this case, the SHG intensity also increased with increasing the concentrations of peptide molecules on the surface. So, the generation of SHG is occurred only from the surface of peptide due to have crystalline directional behavior, no matter whether there is CNT layer or not.